

- Kennedy, R. S., Gonzales, P. C. & Miranda, Jr., H. C. 1997. New *Aethopyga* Sunbirds (Aves: Nectariniidae) from the Island of Mindanao, Philippines. *Auk* 114: 1–10.
- Mearns, E. A. 1906. Description of a new genus and nine new species of Philippine birds. *Phil. J. Sci.* 2: 355–360.
- Meyer de Schauensee, R. & duPont, J. E. 1962. Birds from the Philippine Islands. *Proc. Acad. Nat. Sci. Philadelphia* 114: 149–173.
- Rand, A. L. 1970. Species formation in the blue monarch flycatchers, genus *Hypothymis*. *Nat. Hist. Bull. Siam Soc.* 23: 353–365.
- Ripley, S. D. & Rabor, D. S. 1961. The avifauna of Mt. Katanglad. *Postilla* 50: 1–20.

Addresses: K. R. Reis, Department of Zoology, University of Florida, Gainesville, Florida 32611, U.S.A., email: kreis@zoo.ufl.edu; R. S. Kennedy, Frederick and Amey Geier Collections and Research Center, Museum of Natural History and Science, Cincinnati Museum Center, 1720 Gilbert Avenue, Cincinnati, Ohio 45202, U.S.A.

© British Ornithologists' Club 1999

Comments on the taxonomy of the genus *Cynanthus* (Swainson), with a restricted type locality for *C. doubledayi*

by Adolfo G. Navarro-Sigüenza & A. Townsend Peterson

Received 12 March 1998

Howell & Webb (1995) presented a rearrangement of the taxonomy of the hummingbirds of the Mexican near-endemic genus *Cynanthus*, although without explicit justification for changes proposed. They correctly divided the former *C. latirostris* complex into two portions—that of northern and central Mexico (*C. latirostris*), and that of coastal southwestern Mexico (*C. doubledayi*). In addition, they mentioned the form *C. l. lawrencei* of the Islas Tres Mariás as a distinct subgroup of *C. latirostris*, at times considered a separate species. While we agree with their conclusions, our examination of series of specimens of each form and several of the types leads us to point out a correction necessary for the accurate documentation of the group's taxonomy.

Nominate *latirostris* was described by Swainson (1827) from the "Tableland of Mexico", a locality later corrected to "Valley of Mexico, near Mexico City" by Moore (1939). Subsequent forms described included *doubledayi*, *toroi*, and *nitidus* of Guerrero and southwestern Mexico, *magicus* of northwestern Mexico, *propinquus* of Guanajuato and Michoacán, and *lawrencei* of the Islas Tres Mariás. Clearly, confusion and synonymy are suggested in the plethora of forms named from Guerrero, the subject of the present note.

The name *doubledayi* was first applied by Bourcier (1847) to a specimen apparently now housed in the Loddiges Collection of the Natural History Museum, Tring, allegedly from "Rio Negro", [Brazil]. Subsequently, Salvin & Godman (1889) described *Iache nitida* from a specimen taken by Mrs H. H. Smith at Río Papagaio, Acapulco, carefully contrasting it with what they thought was *doubledayi*:

This beautiful species is most nearly allied with *I. doubledayi*, but the whole throat and breast are of a richer blue, and the head also is shining blue, not green. The specimen of *I. doubledayi* with which we have compared it is stated to be from Chinantla, and is that figured in Gould's 'Monograph of the Humming-Birds'.

Cory (1918) pointed out Bourcier's erroneous type locality, and added a note "=State of Guerrero, Mexico", and a footnote that "the type probably came from State of Guerrero, Mexico". Cory unfortunately did not specify a locality within Guerrero—a source of future confusion. Peters (1945) also cited the correction of the type locality of *doubledayi* as "Guerrero". The synonymy of *nitidus* and *doubledayi* was established by Hartert (1897) and Salvin & Godman (1879–1904), the confusion clearly arising from Salvin and Godman's comparison of the type of *nitidus* with the Gould Chinantla specimen from outside the range of *doubledayi*.

Then began a series of errors and confusions regarding the taxonomy of the group. Ridgway (1892) reported comparisons with "the type of *I. doubledayi*, kindly lent to me by the American Museum of Natural History in New York City". Greenway (1978), however, indicated that the type of *doubledayi* was not at the American Museum—the type of *latirostris*, however, deposited at that institution, carries the name "Bourcier" referring to its authority for the genus *Circe*, and is apparently the source of Ridgway's confusion. A second error was the citation of the subspecies *doubledayi* as ranging in Puebla and northern Oaxaca by Friedmann *et al.* (1950). They also stated erroneously that its type locality had been designated by Gould as Chimantla (=Chiantla, Puebla).

Reviewing specimens of *Cyananthus* in the collections of the U.S. National Museum of Natural History, Natural History Museum (Tring), University of Kansas Natural History Museum, and the Museo de Zoología of the Facultad de Ciencias, Universidad Nacional Autónoma de México, we summarized variation of male plumage characters in the genus (Table 1). This summary of variation, together with comparisons kindly made for us by Michael P. Walters of Bourcier's type of *doubledayi*, indicate that the specimen coincides closely with the populations of coastal southwestern Mexico, differing from the populations of the interior Balsas Basin (*toroi*) in having a blue rather than green forehead and breast. Cory's (1918) designation is correct, but not sufficiently precise—for this reason, we restrict the type locality of *doubledayi* to Acapulco, an important historical collecting locality on the coast of Guerrero.

Considering the geographic patterning of zones of abrupt morphological change, it is possible to divide the genus *Cyananthus* into four diagnosable units. These sets of populations may be considered independent phylogenetic lineages, constituting phylogenetic or evolutionary species, as follows:

1. *C. sordidus* (Gould 1859)—Balsas Depression and surrounding arid highland areas.
2. *C. latirostris* Swainson 1827—Southeast Arizona and northern Mexico south to the plains of Colima and the Balsas Depression,

TABLE 1
Summary of geographic situation, species taxa, and distinguishing characters of named forms of *Cyananthus* hummingbirds

Form	Range	Type locality	Forehead	Throat	Breast and belly	Undertail coverts	Tail cleft
<i>Cyananthus sordidus sordidus</i>	Balsas Depression and Valle de Tehuacán	Oaxaca	Dull brownish green	Dull brown	Dull brown	Dull grey-brown	Moderate
<i>Cyananthus latirostris magicus</i>	Arizona S to Navarrit and Durango	Mazatlán, Sinaloa	Iridescent green	Iridescent blue	Iridescent green	White	Moderate
<i>latirostris propinquus</i>	Central and eastern Mexico El Bajío	Valley of Mexico Irapuato, Guanajuato	Iridescent green	Iridescent blue	Iridescent green	White	Moderate
<i>toroi</i>	Lower Balsas Depression and Plains of Colima	La Estancia, Guerrero	Iridescent green	Iridescent blue	Iridescent green	White	Moderate
<i>Cyananthus doubledayi doubledayi</i>	Coast of southwestern Mexico	Acapulco, Rio Papagayo	Iridescent blue	Iridescent blue	Iridescent blue	Blue-black	Moderate
<i>Cyananthus lawrencei lawrencei</i>	Islas Tres Marias	Islas Tres Marias	Iridescent green	Iridescent green	Iridescent green	White	Deep

- including the present subspecies *latirostris*, *magicus* (Mulsant and Verreaux 1872), *propinquus* Moore 1939, and *toroi* (Berlioz 1937).
3. *C. doubledayi* (Bourcier 1847)—Coastal plain of Mexico in Guerrero, Oaxaca, and extreme southwestern Chiapas.
 4. *C. lawrencei* (Berlepsch 1887)—Confined to the islands María Madre and María Cleofas in the Tres Marias archipelago.

Given the striking differences and apparent parapatry, these phylogenetic/evolutionary species units may also prove to be distinct biological species as well as upon further study. The discrete character variation among *Cynanthus* populations, focused on head and underparts, recalls strikingly the variation in the genus *Thalurania* (Escalante-Pliego and Peterson 1992), although the geographic distributions of the two genera are quite different.

Acknowledgements

We thank James Dean and Gary Graves of the U.S. National Museum of Natural History for assistance and access to collections under their care; Harry D. Lowry, Jr., for his blessing of this effort and for help with logistics; and Michael P. Walters for careful and detailed comments on specimens housed at the Natural History Museum (Tring).

References:

- Bourcier, J. 1847. Description de quinze espèces de Trochilidées du cabinet de M. Loddiges. *Proc. Zool. Soc., Lond.* 15: 42–47.
- Cory, C. B. 1918. *Catalogue of the birds of the Americas and the adjacent islands, Part 2, No. 1.* Field Museum of Natural History, Chicago.
- Escalante P., P. & Peterson, A. T. 1992. Geographic variation and species limits in the hummingbird genus *Thalurania*. *Wilson Bull.* 104: 205–219.
- Friedmann, H., Griscom, L. & Moore, R. T. 1950. Distributional check-list of the birds of Mexico. *Pacific Coast Avifauna* 29: 1–202.
- Greenway, J. C. Jr. 1978. Type specimens of birds in the American Museum of Natural History, Part 2. *Bull. Amer. Mus. Nat. Hist.* 161: 1–305.
- Hartert, E. 1897. Various notes on humming-birds. *Novitates Zool.* 4: 529–533.
- Howell, S. N. G. & Webb, S. 1995. *A guide to the birds of Mexico and northern Central America.* Oxford Univ. Press, Oxford.
- Moore, R. T. 1939. A new race of *Cynanthus latirostris* from Guanajuato. *Proc. Biol. Soc. Washington* 52: 57–60.
- Peters, J. L. 1945. *Check-list of the birds of the world, vol. V.* Harvard Univ. Press, Cambridge, Mass.
- Ridgway, R. 1892. *The hummingbirds.* U.S. National Museum, Washington, D.C.
- Salvin, O. & Godman, F. D. 1879–1904. *Biologia Centrali-Americana.* Taylor & Francis, London.
- Salvin, O. & Godman, F. D. 1889. Notes on Mexican birds. *Ibis* 6th ser., 1: 233–243.
- Swainson, W. 1827. A synopsis of the birds discovered in Mexico by W. Bullock, F.L.S. and H.S. and Mr William Bullock, Jr. *Philosophical Magazine* 1827: 363–442

Address: Adolfo G. Navarro-Sigüenza, Museo de Zoología, Facultad de Ciencias, Universidad Nacional Autónoma de México, Apartado Postal 70-399, México, D.F. 04510 México; A. Townsend Peterson, Natural History Museum and Biodiversity Research Center, The University of Kansas, Lawrence, Kansas 66045.